



## HOT FIREPol<sup>®</sup> DNA Polymerase

(HOT START DNA Polymerase)

Cat. No.	Pack Size	Conc.
01-02-0000S	100 U SAMPLE	5 U/μl
01-02-00500	500 U	5 U/μl
01-02-01000	1000 U	5 U/μl

For *in vitro* use only.

### Description:

HOT FIREPol<sup>®</sup> DNA Polymerase is a chemically modified FIREPol<sup>®</sup> DNA Polymerase. At ambient temperatures it is inactive, having no polymerization activity. HOT FIREPol<sup>®</sup> DNA Polymerase is activated by a 15 min incubation step at 95°C. This prevents extension of non-specifically annealed primers and primer-dimers formed at low temperatures during PCR setup. The enzyme has 5'→3' polymerase and 5' flap endonuclease activity.

### Source:

Purified from an *E. coli* strain that carries an overproducing plasmid containing a modified gene of *Thermus aquaticus* DNA Polymerase.

### Applications:

- Hot Start PCR
- DHPLC
- TA cloning

### Reagents Provided:

- **HOT FIREPol<sup>®</sup> DNA Polymerase**
- **HOT FIREPol<sup>®</sup> 10x Buffer B1** (Mg<sup>2+</sup> and detergent free) *Tris-HCl and (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>*
- **HOT FIREPol<sup>®</sup> 10x Buffer B2** (Mg<sup>2+</sup> free) *Tris-HCl, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and detergent*
- **25 mM MgCl<sub>2</sub>**
- **10x Solution S**  
*Additive that facilitates amplification of difficult templates (e.g. GC-rich DNA templates). This solution should be used at a defined working concentration (1x, 2x or 3x solution).*  
**Solution S is NOT a reaction buffer and should be used ONLY IF non-specific amplifications occur.**

### Concentration:

5 U/μl

### Unit definition:

One unit is defined as the amount of the enzyme required to catalyze the incorporation of 10 nmol of dNTPs into an acid-insoluble form in 30 minutes at 74°C.

### Storage and Dilution buffer:

50% glycerol (v/v), 20 mM Tris-HCl pH 8.7 at 25°C, 100 mM KCl, 0.1 mM EDTA and stabilizers.

### Quality control:

The enzyme is free of nicking and priming activities, exonucleases and non-specific endonucleases. SDS/PAGE - 95 kD band, >98% pure. Activity and stability tested via thermo cycling. The error rate per nucleotide per cycle is ~ 2.5 x 10<sup>-5</sup>; the accuracy is ~ 4 x 10<sup>4</sup>. Estimated half life at 95°C is 1.5 hours.

### Shipping and Storage conditions:

Routine storage: -20°C

Shipping and temporary storage for up to 1 month at room temperature has no detrimental effects on the quality of HOT FIREPol<sup>®</sup> DNA Polymerase.

### Recommended PCR Buffers:

Our standard buffers are:

- **HOT FIREPol<sup>®</sup> 10x Buffer B1** (Mg<sup>2+</sup> and detergent free) *Tris-HCl and (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>*
- **HOT FIREPol<sup>®</sup> 10x Buffer B2** (Mg<sup>2+</sup> free) *Tris-HCl, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and detergent*

For further optimization please try our whole set of buffers available upon request:

- **10x Reaction buffer B** (Mg<sup>2+</sup> free) *0.8 M Tris-HCl, 0.2 M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.2% w/v Tween-20*
- **10x Reaction buffer BD** (Mg<sup>2+</sup> and detergent free) *0.8 M Tris-HCl, 0.2 M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>*

### Recommendations:

Reaction setup at room temperature is highly recommended for HOT FIREPol<sup>®</sup> DNA Polymerase.

In order to prevent contamination, we recommend you to setup the reaction under laminar or in PCR box.

### Recommended PCR reaction mix:

Component	Volume	Final conc.
HOT FIREPol <sup>®</sup> (5 U/μl)	0.4-1.0 μl	0.02-0.05 U/μl (2-5 U)
10x Buffer B1/B2,B/BD	10 μl	1x
25 mM MgCl <sub>2</sub>	6-10 μl	1.5-2.5 mM
dNTP mix (20 mM of each)	1 μl	200 μM of each
Primer Forward (10 pmol/μl)	1-3 μl	0.1-0.3 μM
Primer Reverse (10 pmol/μl)	1-3 μl	0.1-0.3 μM
DNA template <sup>1</sup>	variable <sup>1</sup>	variable <sup>1</sup>
10x Solution S <b>Not for standard PCR</b>	0, 10, 20 or 30 μl	1x, 2x or 3x
H <sub>2</sub> O PCR grade	Up to 100 μl	
<b>Total</b>	<b>100 μl</b>	

<sup>1</sup>Conc. of cDNA 0.01 pg/μl -0.1 ng/μl ; gDNA 0.1 ng/μl – 10 ng/μl

### Recommended PCR cycles:

Cycle step	Temp.	Time	Cycles
<b>Initial activation<sup>2</sup></b>	<b>95°C</b>	<b>12-15 min</b>	1
Denaturation	95°C	15-30 s	26-35
Annealing	50-68°C	30-60 s	
Elongation <sup>3</sup>	72°C	1-4 min	
Final elongation	72°C	5-10 min	1

<sup>2</sup>To activate the polymerase, include an incubation step at **95°C for 12 - 15 minutes** at the beginning of the PCR cycle.

<sup>3</sup>Elongation time should be ~1 min/1 kb.

### Safety warnings and precautions:

This product and its components should be handled only by persons trained in laboratory techniques. It is advisable to wear suitable protective clothing, such as laboratory overalls, gloves and safety glasses. Care should be taken to avoid contact with skin or eyes. In case of contact with skin or eyes, wash immediately with water.

### Related products:

Product name	Pack size	Cat. No.
<b>FIREPol<sup>®</sup> DNA Polymerase</b>	500 U	01-01-00500
<b>FIREPol<sup>®</sup> DNA Polymerase</b>	1000 U	01-01-01000
<b>FIREPol<sup>®</sup> DNA Polymerase</b>	2000 U	01-01-02000
<b>5x FIREPol<sup>®</sup> Master Mix</b> <i>(1.5 mM MgCl<sub>2</sub> final conc.)</i>	250 reactions	04-11-00115
<b>5x FIREPol<sup>®</sup> Master Mix</b> <i>(2.5 mM MgCl<sub>2</sub> final conc.)</i>	250 reactions	04-11-00125
<b>5x FIREPol<sup>®</sup> Master Mix</b> <b>Ready to Load</b> <i>(1.5 mM MgCl<sub>2</sub> final conc.)</i>	250 reactions	04-12-00115
<b>5x FIREPol<sup>®</sup> Master Mix</b> <b>Ready to Load</b> <i>(2.5 mM MgCl<sub>2</sub> final conc.)</i>	250 reactions	04-12-00125
<b>dNTP MIX (20 mM of each)</b>	20 µmol	02-31-00020
<b>dNTP MIX (20 mM of each)</b>	100 µmol	02-31-00100
<b>dNTP SET (100 mM)</b>	4 x 25 µmol	02-21-00100
<b>dNTP SET (100 mM)</b>	4 x 100 µmol	02-21-00400

**Permitted Use:** This product is supplied for research use only (the **Permitted Use**). If the customer wishes to use the product for any purpose other than the Permitted Use, including (without limitation) resale or alteration, the customer should obtain the appropriate licence from Solis BioDyne. Some applications of this product may require a license/licenses from one or more third parties which are not provided by the purchase of this product. Users should obtain the licence if required. Covered by the patent EP2501716, made by the methods of US Patent No 9,321,999.

**Warranty and Disclaimer:** This product shall comply with its relevant specification and be fit for its stated purpose, but Solis BioDyne gives no other warranty and makes no representation as to description or quality. Any such warranty or representation is excluded, to the fullest extent permitted by law. In particular, but without limiting the foregoing, Solis BioDyne shall not be liable for the failure of the product to comply with its relevant specification where such failure arises as a result of: (i) customer negligence or because the customer failed to follow any of the applicable technical data or safety sheets, standard user materials, use guidelines or any other information provided by Solis BioDyne as to the storage, transportation, handling, use or maintenance of the products or other good practice regarding the same, or (ii) the customer altering the products in any way without the prior written consent from Solis BioDyne, or (iii) the products differing from the relevant specification as a result of changes made to ensure their compliance with applicable statutory or regulatory requirements.

Nothing shall limit or exclude Solis BioDyne's liability for death or personal injury caused by its negligence, fraud or fraudulent misrepresentation or any matter in respect of which it would be unlawful for Solis BioDyne to exclude or restrict liability. Without limiting the foregoing, Solis BioDyne shall under no circumstances whatever be liable to the customer, whether in contract, tort (including negligence), breach of statutory duty, or otherwise, for any loss of profit, or any indirect or consequential loss arising under or in connection with the products and Solis BioDyne's total liability to the customer in respect of all other losses arising under or in connection with the product, whether in contract, tort (including negligence), breach of statutory duty, or otherwise, shall in no circumstances exceed the price of the products supplied in respect of which the liability has arisen.